

Rockefeller University

Digital Commons @ RU

---

Publications

Steinman Laboratory Archive

---

2000

## Will the making of plasmacytoid dendritic cells in vitro help unravel their mysteries?

Jacques Banchereau

Bali Pulendran

Ralph M. Steinman

Karolina A. Palucka

Follow this and additional works at: <https://digitalcommons.rockefeller.edu/steinman-publications>

---

## Will the Making of Plasmacytoid Dendritic Cells In Vitro Help Unravel Their Mysteries?

By Jacques Banchereau,\* Bali Pulendran,\* Ralph Steinman,‡  
and Karolina Palucka\*

From the \*Baylor Institute for Immunology Research, Dallas, Texas 75204; and the ‡Laboratory of Cellular Physiology and Immunology, The Rockefeller University, New York, New York 10021

In this issue, two papers describe the in vitro generation of an intriguing subset of human dendritic cells (DCs), the plasmacytoid DCs (pDCs [1, 2]). For nearly 20 years after their discovery (3), DCs had to be painstakingly isolated from tissues (4). Progress was rather slow, as DCs represent a minor cell population in all tissues, and relatively few investigators appreciated their importance in antigen presentation and the control of immunity. In 1992, in vitro culture systems were identified to generate large numbers of mouse (5) and human DCs (6). These culture systems considerably accelerated the study of DCs, with many groups joining the search to unravel their mysteries (7–9). Currently, DCs are being developed in vitro from (a) bone marrow progenitors cultured in GM-CSF, with TNF being essential for human cells (6), and (b) human blood monocytes cultured with GM-CSF and IL-4 (10–12). Recently, Flt3 ligand (Flt3-L), a stromal cell product, was found to induce a massive expansion of DCs in vivo in mice (13, 14) and in humans (15, 16).

**Subsets of DCs.** The concept of distinct DC subsets in humans came from several routes including analyses of (a) skin DCs (17), (b) DCs generated in vitro by culture of CD34<sup>+</sup> hematopoietic progenitors (HPCs [18]), and (c) blood DC precursors (19). Human skin contains two distinct subsets: Langerhans cells (LCs) within the epidermis, characterized by the expression of CD1a and Birbeck granules, and interstitial (dermal) DCs, lacking Birbeck granules but expressing coagulation factor XIIIa. These two subsets also emerge in cultures of CD34<sup>+</sup> HPCs driven by the addition of GM-CSF plus TNF- $\alpha$  (18). These subsets have common as well as unique functions (20). In particular, interstitial DCs, but not LCs, are able to induce the differentiation of naive B cells into immunoglobulin-secreting plasma cells. Although no unique function has yet been formally attributed to LCs, there are hints they may be particularly efficient activators of cytotoxic CD8 T cells (21, 22). Two subsets of DCs were identified in the blood, each representing a small fraction ( $\sim 0.3\%$ ) of the mononuclear

cells (19). One subset, CD11c<sup>+</sup> population, differentiated into mature DCs in response to inflammatory stimuli, whereas a second CD11c<sup>-</sup> subset was prone to prompt apoptosis in culture, and was later identified as “pDCs.”

**pDCs and “DC2.”** Plasmacytoid cells were first observed by experts in the histopathology of lymphoid organs (23–26) who saw cells within the T cell areas that looked like plasma cells. Because plasma cells are normally localized outside the standard B cell follicles and T cell areas, it was concluded that these “plasmacytoid” cells were different. Indeed, due to the expression of CD68 (27) and CD4, these cells were sometimes called either plasmacytoid monocytes or plasmacytoid T cells. Their isolation from tonsils by Grouard et al. revealed a unique phenotype (28). Transmission electron microscopy showed an abundant rough endoplasmic reticulum and an eccentric nucleus, quite similar to plasma cells. Flow cytometry demonstrated the expression of IL-3 receptor, conferring responsiveness to IL-3 but sparse expression of GM-CSF receptor, consistent with the lack of responsiveness to GM-CSF (29). In addition, many other myeloid markers (such as CD13, CD14, and CD33) were missing. pDCs can be further distinguished from CD11c<sup>+</sup> DCs by differential expression of immunoglobulin-like transcripts (ILTs), with pDCs being ILT1<sup>-</sup>ILT3<sup>+</sup> and CD11c<sup>+</sup> DCs being ILT1<sup>+</sup>ILT3<sup>+</sup> (30). The latter report emphasized that pDCs home to the T cell areas of lymphoid tissues, e.g., via L-selectin. At the International Workshop on DCs held in Pittsburgh in September 1998, Liu and co-workers presented two novel exciting properties of these cells: their ability to produce large amounts of IFN- $\alpha$  and their ability to polarize a fraction of T cells towards IL-4 and IL-5 production (type 2 cells [31]).

This rare cell type has been given many names, including pDCs, CD11c<sup>-</sup> DCs, DC2, and IFN-producing cells (IPCs). For the sake of simplicity, and by reference to the terminology used in the two papers from the current issue, we shall call these cells “DC2.” It is anticipated that two recent findings will accelerate research on human DC2. First, systemic administration of both Flt3-L and G-CSF increase the number of DC2 in blood (15, 16, 32), and second, the new studies presented in this issue provide culture systems to generate DC2 in vitro. The group of Blom et al.

Address correspondence to J. Banchereau, Baylor Institute for Immunology Research, 3434 Live Oak, Dallas, TX 75204. Phone: 214-820-7450; Fax: 214-820-4813; E-mail: j.banchereau@bayloridallas.edu

(1) demonstrates the existence of CD34<sup>low</sup> cells with the phenotype and function of DC2, as well as a simple culture system in which early CD34<sup>+</sup> progenitors yield DC2 (1). The group of Spits et al. (2) provides another culture system in which early CD34<sup>+</sup> progenitors yield DC2, and demonstrates that their differentiation can be blocked by overexpression of inhibitor of DNA binding (Id)2 and Id3 proteins (2).

**CD34<sup>+</sup> Cells with Features of DC2.** Several features permit the identification of DC2s: the expression of CD4, CD45RA, and CD123, the production of high levels of IFN- $\alpha$  upon viral triggering or CD40 ligation (30, 33, 34), and the morphology of plasma cells. Blom et al. analyzed hematopoietic tissues and looked for the presence of CD34<sup>+</sup> cells with the features of DC2 (1). Accordingly, a minor subset of CD34<sup>low</sup>CD45RA<sup>+</sup>CD4<sup>+</sup>CD123<sup>+</sup> cells was identified in cord blood, adult blood, fetal liver, and most abundantly, fetal bone marrow. These cells produce large amounts of IFN- $\alpha$  in response to viruses and can differentiate into mature DCs in response to IL-3 and CD40-L.

**In Vitro Generation of DC2 from Early CD34<sup>+</sup> Hematopoietic Progenitors.** Culturing early HPCs (CD34<sup>+</sup>CD45RA<sup>-</sup>) with Flt3-L gives rise to cells (a) able to secrete IFN- $\alpha$  upon viral exposure and (b) displaying the typical DC2 phenotype (CD4<sup>+</sup>CD123<sup>+</sup>CD45RA<sup>+</sup>). The generation of DC2 under these conditions is slow, as they are first observed after 10 d and reach a maximal frequency ( $\sim$ 10%) after 3 wk of culture. The identification of Flt3-L as the DC2 hematopoietic differentiation factor does not come as a surprise inasmuch as Flt3-L increases circulating DC2. More surprising is the fact that G-CSF, which also increases DC2 in vivo, does not permit the generation of these cells in vitro. Thus, G-CSF acts in vivo as a mobilizing agent for DC2, as it does for CD34<sup>+</sup> HPCs (35).

Using a different approach, Spits et al. were able to generate DC2 by culturing early uncommitted CD34<sup>+</sup>CD38<sup>-</sup> fetal liver HPCs on the murine stromal cell line S17 (2). This cell line is unusual in that it supports the development of human B cell and myeloid progenitors (36–38). At variance with the above system, it takes only a few days to see the emergence of DC2, but then the cells quickly disappear. The newly generated cells express pre-TCR- $\alpha$  transcripts, unlike the starting CD34<sup>+</sup> HPCs. They also display phenotypic and functional features of DC2: they secrete IFN- $\alpha$  upon viral challenge and differentiate into mature DCs after exposure to IL-3 and CD40-L. Cells with the same properties can be obtained from thymic CD34<sup>+</sup> progenitors, but from the CD34<sup>+</sup>CD1a<sup>-</sup> subset only. Thus, the cells generated in Spits et al. culture system (2) appear similar to those obtained in Blom et al. (1) system.

Because the kinetics of DC2 growth and survival are so different in the two culture systems, we surmise that the S17 system may reveal a DC2 growth factor different from Flt3-L.


**Arguments for the Lymphoid Origin of DC2.** The term “lymphoid DC” was coined by Shortman and colleagues to describe a mouse DC subset in the thymus that developed from a population of thymic lymphoid progenitor cells in

vivo (39). In these studies, transfer of purified thymic lymphoid precursors into irradiated hosts resulted in the development of T cells, B cells, NK cells, and DCs expressing CD8 $\alpha$ <sup>+</sup>, but not cells of the myeloid lineage (39, 40). These were striking findings because previous systems for developing DCs also led to the production of macrophages and granulocytes. Based on these observations, CD8 $\alpha$  was postulated to be a characteristic marker for lymphoid DCs in mice. Particularly intriguing was the postulate that lymphoid DCs play a role in the induction of central tolerance (41, 42). CD8 $\alpha$ <sup>+</sup> DCs with a similar phenotype to thymic DCs were also observed in the spleens, lymph nodes, and Peyer’s patch, and these DCs have been presumed to be of lymphoid origin as well (43). In this context, the role of DCs in the establishment and maintenance of peripheral tolerance has recently been discussed (44, 45).

It should be noted that there is no clonal analysis to support the claim that CD8 $\alpha$ <sup>+</sup> DCs and lymphoid cells arise from the same precursor cell. Furthermore, recent studies suggest that CD8 $\alpha$  can also be expressed on activated DCs and LCs (46, 47), a cell type that has hitherto been considered to be of myeloid origin. Therefore, at present the lineage origins of DC subsets in mice are a matter of considerable debate.

The concept of a lymphoid DC subset in mice sparked great interest in the search for lymphoid DCs in humans. Unlike in mice, human thymic DCs do not express CD8, but do express CD4 (48), rendering the phenotypic discrimination between human thymic DCs and myeloid DCs difficult. However, the identification of DC2 and some of their features has offered new clues. Despite their lack of expression of CD11c and CD8a (two markers expressed by murine lymphoid DCs), several features of human DC2 are consistent with a lymphoid origin: (a) DC2 lack expression of the myeloid antigens CD11c, CD13, and CD33; (b) DC2 isolated from the thymus, express the lymphoid markers CD2, CD5, and CD7; (c) DC2 do not differentiate into macrophages after culture with GM-CSF and M-CSF; and (d) the thymic DC precursors as well as the tonsillar DC2 expressed significant levels of pre-TCR- $\alpha$  transcripts (49). Spits et al. now provide another element to this claim using the S17 system described above (2). Transduction of early CD34 progenitors with Id2 and Id3 (proteins that prevent the binding of basic helix-loop-helix transcription factors to DNA) inhibits their differentiation into DC2 as well as T cells. In contrast, generation of CD4<sup>+</sup>CD14<sup>+</sup> cells, as well as the development of DCs in a standard culture system of CD34<sup>+</sup> HPCs with GM-CSF and TNF, are not affected. Furthermore, Id3 inhibits the development of primitive HPCs into T cells and B cells while stimulating that of NK cells (50). These observations therefore support the notion of shared molecular cues for the development of DC2 precursors, T and B cells (Fig. 1).

**DCs: Regulators of Innate and Adaptive Immunity.** Although DCs have long been recognized to initiate the immune response, their roles in regulating both the innate and adaptive arms of immunity are beginning to be elucidated. DCs have a role in innate immunity in that they



**Figure 1.** A summary of potential differentiation pathways of human DCs. CD34<sup>+</sup> myeloid progenitors give rise to monocytes, a large reservoir of antigen presenting cell precursors that yield circulating CD11c<sup>+</sup> precursors. CD11c<sup>+</sup> cells can differentiate into either macrophages or interstitial DCs or LCs depending on the cytokine microenvironment (reference 64). A distinct precursor population may give rise to plasmacytoid DCs. In some circumstances, this population also contains cells that can yield lymphocytes, hence the current term “lymphoid.” CD34<sup>+</sup> cells contain a subset with the phenotypic and functional characteristics of DC2, pro-DC2, with a limited proliferative capacity. Thus, human blood contains two DC precursors, CD11c<sup>+</sup> (myeloid) and CD11c<sup>-</sup> (potentially lymphoid). Two main questions need to be resolved regarding these putative lymphoid DCs. There is no clonal system to produce plasmacytoid DCs and T cells or B cells. The classical immature (antigen capturing) and mature (antigen presenting) DC differentiation stages have not yet been characterized, in contrast to myeloid DCs derived from skin, blood, and bone marrow in culture.

have nonclonotypic receptors that permit a prompt response to microbial antigens, leading to production of cytokines that mobilize NK cells and have antimicrobial activity. For example, human monocyte-derived DCs certainly make abundant amounts of IL-12 (51, 52), whereas DC2 make IFN- $\alpha$  (34, 53).

DCs also have a powerful role in regulating the adaptive immune responses, as distinct DC subsets induce different Th responses. In mice, there is a consensus that CD8 $\alpha$ <sup>+</sup> (putative lymphoid) DCs prime Th1 responses, whereas CD8 $\alpha$ <sup>-</sup> (putative myeloid) DCs prime Th2 responses (54, 55). In humans, the picture is less clear. Monocyte-derived DCs activated with CD40-L have been shown to prime Th1 responses through an IL-12-dependent mechanism, whereas IL-3<sup>+</sup> CD40-L-activated DC2s have been shown to secrete negligible amounts of IL-12 and can prime Th2 responses. However, later it was observed that the polarizing effects of DC subsets was not immutable, as different environmental cues could instruct a given DC subset to elicit different Th responses. Thus, monocyte-derived DCs can induce T cells to make IL-4 if (a) the DCs are used at low numbers (56), (b) after exposure to factors such as prostaglandin E2 (PGE2), corticosteroids, or IL-10, or (c)

upon prolonged activation with CD40-L (>3 d in vitro culture; references 57–60). Furthermore, DC2s, when stimulated by virus (rather than IL-3<sup>+</sup> CD40-L), secrete IFN- $\alpha$  that drives Th1 responses in humans (61, 62), and mature into DCs that can induce T cells to produce IFN- $\gamma$  and IL-10 (34, 53). Thus, both the type of DC subset as well as microenvironmental cues are important for Th polarization. Another recently proposed parameter is the stage (early or late) of DCs activation. According to this “DC exhaustion model,” monocyte-derived DCs produce IL-12 shortly after stimulation with lipopolysaccharide, thereby inducing Th1 differentiation. After the burst of IL-12 production, DCs no longer polarize T cells into the Th1 pathway but into the Th2 pathway (9, 63).

To summarize the current literature, there are three models through which DCs may control T cell polarization: (a) subset of DCs; (b) the type of DC maturation or activation signals; and (c) the duration of DC activation. It would not surprise us if each operates under different circumstances of the immune response.

**Conclusions.** DC2 have several striking features: their distinctive morphology and surface markers, their localization in the T cell areas in vivo, their responsiveness to IL-3

and not to GM-CSF, their capacity to induce T cells to make either IFN- $\gamma$  or IL-4 under different conditions, and their prodigious production of IFN- $\alpha$  upon challenge with many enveloped viruses. Two papers in this issue provide methodology to make these cells in vitro. This will help investigators learn more about their biology and to exploit them for therapeutic purposes. The demonstration of the critical role for Flt3-L and stromal cells in the generation of these cells, and their unique ability to secrete large amounts of IFN- $\alpha$  may permit the identification of the mouse equivalent. This should lead to identification of mouse models for preclinical studies aimed at establishing DC2 as targets and vectors for therapy.

Submitted: 8 November 2000

Accepted: 15 November 2000

## References

- Blom, B., S. Ho, S. Antonenko, and Y.J. Liu. 2000. Generation of interferon  $\alpha$ -producing pre-dendritic cell (DC)2 from human CD34<sup>+</sup> hematopoietic stem cells. *J. Exp. Med.* 192: 1785–1795.
- Spits, H., F. Couwenberg, A.Q. Bakker, K. Weijer, and C.H. Uittenbogaart. 2000. Id2 and Id3 inhibit development of CD34<sup>+</sup> stem cells into predendritic cell (pre-DC)2 but not into pre-DC1: evidence for a lymphoid origin of pre-DC2. *J. Exp. Med.* 192:1775–1783.
- Steinman, R.M., and Z.A. Cohn. 1973. Identification of a novel cell type in peripheral lymphoid organs of mice. I. Morphology, quantitation, tissue distribution. *J. Exp. Med.* 137:1142–1162.
- Steinman, R.M. 1991. The dendritic cell system and its role in immunogenicity. *Annu. Rev. Immunol.* 9:271–296.
- Inaba, K., M. Inaba, N. Romani, H. Aya, M. Deguchi, S. Ikehara, S. Muramatsu, and R.M. Steinman. 1992. Generation of large numbers of dendritic cells from mouse bone marrow cultures supplemented with granulocyte/macrophage colony-stimulating factor. *J. Exp. Med.* 176:1693–1702.
- Caux, C., C. Dezutter-Dambuyant, D. Schmitt, and J. Banchereau. 1992. GM-CSF and TNF- $\alpha$  cooperate in the generation of dendritic Langerhans cells. *Nature.* 360: 258–261.
- Banchereau, J., and R.M. Steinman. 1998. Dendritic cells and the control of immunity. *Nature.* 392:245–252.
- Banchereau, J., F. Briere, C. Caux, J. Davoust, S. Lebecque, Y. Liu, B. Pulendran, and K. Palucka. 2000. Immunobiology of dendritic cells. *Annu. Rev. Immunol.* 18:767–811.
- Lanzavecchia, A., and F. Sallusto. 2000. Dynamics of T lymphocyte responses: intermediates, effectors, and memory cells. *Science.* 290:92–97.
- Peters, J.H., H. Xu, J. Ruppert, D. Ostermeier, D. Friedrichs, and R.K. Gieseler. 1993. Signals required for differentiating dendritic cells from human monocytes in vitro. *Adv. Exp. Med. Biol.* 329:275–280.
- Romani, N., S. Gruner, D. Brang, E. Kampgen, A. Lenz, B. Trockenbacher, G. Konwalinka, P.O. Fritsch, R.M. Steinman, and G. Schuler. 1994. Proliferating dendritic cell progenitors in human blood. *J. Exp. Med.* 180:83–93.
- Sallusto, F., and A. Lanzavecchia. 1994. Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor plus interleukin 4 and downregulated by tumor necrosis factor  $\alpha$ . *J. Exp. Med.* 179:1109–1118.
- Maraskovsky, E., K. Brasel, M. Teepe, E.R. Roux, S.D. Lyman, K. Shortman, and H.J. McKenna. 1996. Dramatic increase in the numbers of functionally mature dendritic cells in Flt3 ligand-treated mice: multiple dendritic cell subpopulations identified. *J. Exp. Med.* 184:1953–1962.
- Pulendran, B., J. Lingappa, M.K. Kennedy, J. Smith, M. Teepe, A. Rudensky, C.R. Maliszewski, and E. Maraskovsky. 1997. Developmental pathways of dendritic cells in vivo: distinct function, phenotype, and localization of dendritic cell subsets in FLT3 ligand-treated mice. *J. Immunol.* 159:2222–2231.
- Maraskovsky, E., E. Daro, E. Roux, M. Teepe, C.R. Maliszewski, J. Hoek, D. Caron, M.E. Lebsack, and H.J. McKenna. 2000. In vivo generation of human dendritic cell subsets by Flt3 ligand. *Blood.* 96:878–884.
- Pulendran, B., J. Banchereau, S. Burkeholder, E. Kraus, E. Guinet, C. Chalouni, D. Caron, C. Maliszewski, J. Davoust, J. Fay, and K. Palucka. 2000. Flt3-ligand and granulocyte colony-stimulating factor mobilize distinct human dendritic cell subsets in vivo. *J. Immunol.* 165:566–572.
- Cerio, R., C.E. Griffiths, K.D. Cooper, B.J. Nickoloff, and J.T. Headington. 1989. Characterization of factor XIIIa positive dermal dendritic cells in normal and inflamed skin. *Br. J. Dermatol.* 121:421–431.
- Caux, C., B. Vanbervliet, C. Massacrier, C. Dezutter-Dambuyant, B. de Saint-Vis, C. Jacquet, K. Yoneda, S. Imamura, D. Schmitt, and J. Banchereau. 1996. CD34<sup>+</sup> hematopoietic progenitors from human cord blood differentiate along two independent dendritic cell pathways in response to GM-CSF+TNF  $\alpha$ . *J. Exp. Med.* 184:695–706.
- O'Doherty, U., M. Peng, S. Gezelter, W.J. Swiggard, M. Betjes, N. Bhardwaj, and R.M. Steinman. 1994. Human blood contains two subsets of dendritic cells, one immunologically mature and the other immature. *Immunology.* 82: 487–493.
- Caux, C., C. Massacrier, B. Vanbervliet, B. Dubois, I. Durand, M. Cella, A. Lanzavecchia, and J. Banchereau. 1997. CD34<sup>+</sup> hematopoietic progenitors from human cord blood differentiate along two independent dendritic cell pathways in response to granulocyte-macrophage colony-stimulating factor plus tumor necrosis factor  $\alpha$ : II. Functional analysis. *Blood.* 90:1458–1470.
- Mortarini, R., A. Anichini, M. Di Nicola, S. Siena, M. Bregni, F. Belli, A. Molla, A.M. Gianni, and G. Parmiani. 1997. Autologous dendritic cells derived from CD34<sup>+</sup> progenitors and from monocytes are not functionally equivalent antigen-presenting cells in the induction of melan-A/Mart-1(27–35)-specific CTLs from peripheral blood lymphocytes of melanoma patients with low frequency of CTL precursors. *Cancer Res.* 57:5534–5541.
- Ferlazzo, G., A. Wesa, W.Z. Wei, and A. Galy. 1999. Dendritic cells generated either from CD34<sup>+</sup> progenitor cells or from monocytes differ in their ability to activate antigen-specific CD8<sup>+</sup> T cells. *J. Immunol.* 163:3597–3604.
- Muller-Hermelink, H., H. Stein, G. Steinman, and K. Lennert. 1983. Malignant lymphoma of plasmacytoid T cells. Morphologic and immunologic studies characterizing a special type of T cell. *Am. J. Surg. Pathol.* 8:849–862.
- Vollenweider, R., and K. Lennert. 1983. Plasmacytoid T-cell clusters in non-specific lymphadenitis. *Virchows Arch. B Cell Pathol. Incl. Mol. Pathol.* 44:1–14.

25. Müller-Hermelink, H.K., E. Kaiserling, and K. Lennert. 1973. [Pseudofollicular nests of plasmacells (of a special type?) in paracortical pulp of human lymph nodes]. *Virchows Arch. B Cell Pathol.* 14:47–56.
26. Lennert, K., E. Kaiserling, and H.K. Müller-Hermelink. 1975. Letter: T-associated plasma-cells. *Lancet* 1:1031–1032.
27. Strobl, H., C. Scheinecker, E. Riedl, B. Csmarits, C. Bello-Fernandez, W.F. Pickl, O. Majdic, and W. Knapp. 1998. Identification of CD68<sup>+</sup>lin<sup>-</sup> peripheral blood cells with dendritic precursor characteristics. *J. Immunol.* 161:740–748.
28. Grouard, G., M.C. Rissoan, L. Filgueira, I. Durand, J. Banchereau, and Y.J. Liu. 1997. The enigmatic plasmacytoid T cells develop into dendritic cells with interleukin (IL)-3 and CD40-ligand. *J. Exp. Med.* 185:1101–1111.
29. Olweus, J., A. BitMansour, R. Warnke, P.A. Thompson, J. Carballido, L.J. Picker, and F. Lund-Johansen. 1997. Dendritic cell ontogeny: a human dendritic cell lineage of myeloid origin. *Proc. Natl. Acad. Sci. USA.* 94:12551–12556.
30. Cella, M., D. Jarrossay, F. Facchetti, O. Alebardi, H. Nakajima, A. Lanzavecchia, and M. Colonna. 1999. Plasmacytoid monocytes migrate to inflamed lymph nodes and produce high levels of type I IFN. *Nat. Med.* 5:919–923.
31. Siegal, F.P., N. Kadowaki, M. Shodell, and Y.J. Liu. 1998. Evidence for identity of the natural interferon producing cells (NIPC) and the CD11c<sup>-</sup> precursor of DC2. *J. Leukoc. Biol. Suppl.* 2:18.
32. Arpinati, M., C.L. Green, S. Heimfeld, J.E. Heuser, and C. Anasetti. 2000. Granulocyte-colony stimulating factor mobilizes T helper 2-inducing dendritic cells. *Blood.* 95:2484–2490.
33. Siegal, F.P., N. Kadowaki, M. Shodell, P.A. Fitzgerald-Bocarsly, K. Shah, S. Ho, S. Antonenko, and Y.J. Liu. 1999. The nature of the principal type 1 interferon-producing cells in human blood. *Science.* 284:1835–1837.
34. Kadowaki, N., S. Antonenko, J.Y. Lau, and Y.J. Liu. 2000. Natural interferon  $\alpha/\beta$ -producing cells link innate and adaptive immunity. *J. Exp. Med.* 192:219–226.
35. Bensinger, W.I., R.A. Clift, C. Anasetti, F.A. Appelbaum, T. Demirer, S. Rowley, B.M. Sandmaier, B. Torok-Storb, R. Storb, and C.D. Buckner. 1996. Transplantation of allogeneic peripheral blood stem cells mobilized by recombinant human granulocyte colony stimulating factor. *Stem Cells.* 14: 90–105.
36. Collins, L.S., and K. Dorshkind. 1987. A stromal cell line from myeloid long-term bone marrow cultures can support myelopoiesis and B lymphopoiesis. *J. Immunol.* 138:1082–1087.
37. Rawlings, D.J., S. Quan, Q.L. Hao, F.T. Thiemann, M. Smogorzewska, O.N. Witte, and G.M. Crooks. 1997. Differentiation of human CD34<sup>+</sup>CD38<sup>-</sup> cord blood stem cells into B cell progenitors in vitro. *Exp. Hematol.* 25:66–72.
38. Hao, Q.L., E.M. Smogorzewska, L.W. Barsky, and G.M. Crooks. 1998. In vitro identification of single CD34<sup>+</sup>CD38<sup>-</sup> cells with both lymphoid and myeloid potential. *Blood.* 91: 4145–4151.
39. Ardavin, C., L. Wu, C.L. Li, and K. Shortman. 1993. Thymic dendritic cells and T cells develop simultaneously in the thymus from a common precursor population. *Nature.* 362: 761–763.
40. Wu, L., C.L. Li, and K. Shortman. 1996. Thymic dendritic cell precursors: relationship to the T lymphocyte lineage and phenotype of the dendritic cell progeny. *J. Exp. Med.* 184: 903–911.
41. Matzinger, P., and S. Guerder. 1989. Does T-cell tolerance require a dedicated antigen-presenting cell? *Nature.* 338:74–76.
42. Shortman, K., D. Vremec, L.M. Corcoran, K. Georgopoulos, K. Lucas, and L. Wu. 1998. The linkage between T-cell and dendritic cell development in the mouse thymus. *Immunol. Rev.* 165:39–46.
43. Shortman, K. 2000. Burnet oration: dendritic cells: multiple subtypes, multiple origins, multiple functions. *Immunol. Cell Biol.* 78:161–165.
44. Steinman, R.M., M. Pack, and K. Inaba. 1997. Dendritic cells in the T-cell areas of lymphoid organs. *Immunol. Rev.* 156:25–37.
45. Steinman, R.M., S. Turley, I. Mellman, and K. Inaba. 2000. The induction of tolerance by dendritic cells that have captured apoptotic cells. *J. Exp. Med.* 191:411–416.
46. Merad, M., L. Fong, J. Bogenberger, and E.G. Engleman. 2000. Differentiation of myeloid dendritic cells into CD8 $\alpha$ -positive dendritic cells in vivo. *Blood.* 96:1865–1872.
47. Anjuere, F., G.M. del Hoyo, P. Martin, and C. Ardavin. 2000. Langerhans cells develop from a lymphoid-committed precursor. *Blood.* 96:1633–1637.
48. Sotzik, F., Y. Rosenberg, A.W. Boyd, M. Honeyman, D. Metcalf, R. Scollay, L. Wu, and K. Shortman. 1994. Assessment of CD4 expression by early T precursor cells and by dendritic cells in the human thymus. *J. Immunol.* 152:3370–3377.
49. Res, P.C., F. Couwenberg, F.A. Vyth-Dreese, and H. Spits. 1999. Expression of pT $\alpha$  mRNA in a committed dendritic cell precursor in the human thymus. *Blood.* 94:2647–2657.
50. Heemskerk, M.H., B. Blom, G. Nolan, A.P. Stegmann, A.Q. Bakker, K. Weijer, P.C. Res, and H. Spits. 1997. Inhibition of T cell and promotion of natural killer cell development by the dominant negative helix loop helix factor Id3. *J. Exp. Med.* 186:1597–1602.
51. Heufler, C., F. Koch, U. Stanzl, G. Topar, M. Wysocka, G. Trinchieri, A. Enk, R.M. Steinman, N. Romani, and G. Schuler. 1996. Interleukin-12 is produced by dendritic cells and mediates T helper 1 development as well as interferon-gamma production by T helper 1 cells. *Eur. J. Immunol.* 26: 659–668.
52. Cella, M., D. Scheidegger, K. Palmer-Lehmann, P. Lane, A. Lanzavecchia, and G. Alber. 1996. Ligation of CD40 on dendritic cells triggers production of high levels of interleukin-12 and enhances T cell stimulatory capacity: T-T help via APC activation. *J. Exp. Med.* 184:747–752.
53. Cella, M., F. Facchetti, A. Lanzavecchia, and M. Colonna. 2000. Plasmacytoid dendritic cells activated by influenza virus and CD40 ligand drive a potent Th1 polarization. *Nat. Immunol.* 1:305–310.
54. Pulendran, B., J.L. Smith, G. Caspary, K. Brasel, D. Pettit, E. Maraskovsky, and C.R. Maliszewski. 1999. Distinct dendritic cell subsets differentially regulate the class of immune response in vivo. *Proc. Natl. Acad. Sci. USA.* 96:1036–1041.
55. Maldonado-Lopez, R., T. De Smedt, P. Michel, J. Godfroid, B. Pajak, C. Heirman, K. Thielemans, O. Leo, J. Urbain, and M. Moser. 1999. CD8 $\alpha$ <sup>+</sup> and CD8 $\alpha$ <sup>-</sup> subclasses of dendritic cells direct the development of distinct T helper cells in vivo. *J. Exp. Med.* 189:587–592.
56. Tanaka, H., C.E. Demeure, M. Rubio, G. Delespesse, and M. Sarfati. 2000. Human monocyte-derived dendritic cells

- induce naive T cell differentiation into T helper cell type 2 (Th2) or Th1/Th2 effectors. Role of stimulator/responder ratio. *J. Exp. Med.* 192:405–412.
57. Vieira, P.L., P. Kalinski, E.A. Wierenga, M.L. Kapsenberg, and E.C. de Jong. 1998. Glucocorticoids inhibit bioactive IL-12p70 production by in vitro-generated human dendritic cells without affecting their T cell stimulatory potential. *J. Immunol.* 161:5245–5251.
  58. Kalinski, P., J.H. Schuitemaker, C.M. Hilkens, and M.L. Kapsenberg. 1998. Prostaglandin E2 induces the final maturation of IL-12-deficient CD1a<sup>+</sup>CD83<sup>+</sup> dendritic cells: the levels of IL-12 are determined during the final dendritic cell maturation and are resistant to further modulation. *J. Immunol.* 161:2804–2809.
  59. Kalinski, P., J.H. Schuitemaker, C.M. Hilkens, E.A. Wierenga, and M.L. Kapsenberg. 1999. Final maturation of dendritic cells is associated with impaired responsiveness to IFN-gamma and to bacterial IL-12 inducers: decreased ability of mature dendritic cells to produce IL-12 during the interaction with Th cells. *J. Immunol.* 162:3231–3236.
  60. Vieira, P.L., E.C. de Jong, E.A. Wierenga, M.L. Kapsenberg, and P. Kalinski. 2000. Development of Th1-inducing capacity in myeloid dendritic cells requires environmental instruction. *J. Immunol.* 164:4507–4512.
  61. Parronchi, P., S. Mohapatra, S. Sampognaro, L. Giannarini, U. Wahn, P. Chong, E. Maggi, H. Renz, and S. Romagnani. 1996. Effects of interferon-alpha on cytokine profile, T cell receptor repertoire and peptide reactivity of human allergen-specific T cells. *Eur. J. Immunol.* 26:697–703.
  62. Rogge, L., L. Barberis-Maino, M. Biffi, N. Passini, D.H. Presky, U. Gubler, and F. Sinigaglia. 1997. Selective expression of an interleukin-12 receptor component by human T helper 1 cells. *J. Exp. Med.* 185:825–831.
  63. Langenkamp, A., M. Messi, A. Lanzavecchia, and F. Sallusto. 2000. Kinetics of dendritic cell activation: impact on priming of TH1, TH2 and nonpolarized T cells. *Nat. Immunol.* 1:311–316.
  64. Ito, T., M. Inaba, K. Inaba, J. Toki, S. Sogo, T. Iguchi, Y. Adachi, K. Yamaguchi, R. Amakawa, J. Valladeau, et al. 1999. A CD1a<sup>+</sup>/CD11c<sup>+</sup> subset of human blood dendritic cells is a direct precursor of Langerhans cells. *J. Immunol.* 163:1409–1419.